ECOLOGY, BEHAVIOR AND BIONOMICS



	ENTC	MOLO	^{ÓGI} CA	
JOHO		20/	\bigcirc	00
		26		8 P A B
	- <u></u> ,	1972	• 71	

Is Planting Trees Enough? The Effect of Different Types of Reforestation on the Offspring of *Trypoxylon (Trypargilum) lactitarse* (Hymenoptera: Crabronidae) in the Southern Amazon

GJ de Araújo¹, D Storck-Tonon², TJ Izzo¹

¹Dept de Ecologia e Botânica, Lab de Ecologia de Comunidades, Univ Federal de Mato Grosso, Cuiabá, Mato Grosso, Brasil ²Programa de Pós-Graduação em Ambiente e Sistemas de Produção Agrícola, Univ do Estado de Mato Grosso, Cáceres, Brasil

Keywords

Environmental recovery, ecological succession, tropical forests, population patterns, insect bioindicators

Correspondence

Gustavo Júnior de Araújo, Dept de Ecologia e Botânica, Lab de Ecologia de Comunidades, Univ Federal de Mato Grosso, Cuiabá, Mato Grosso, Brasil; gustavojraraujo@gmail.com

Edited by Heraldo Vasconcelos - UFU

Received 21 September 2018 and accepted 17 March 2019 Published online: 9 April 2019

© Sociedade Entomológica do Brasil 2019

Introduction

In the last decades, the increase in agriculture and cattle ranching activities has been the main cause of the disappearance of tropical forests (Fearnside 2006, Newbold *et al* 2015). In the Amazon rainforest, thousands of hectares are lost every year and, as a consequence, many species of the flora and fauna are extinct (Gibbs *et al* 2010, Laurance *et al* 2011). Associated to these extinctions, there is the loss of

Abstract

The deforestation has led to local loss of species and important ecosystem services performed by them, causing ecological and economic losses. It is proposed that the reforestation of such areas aims to reduce those impacts. However, particularly in the tropics, little is known about the real success of different types of reforestation in the recovery of the species, and especially of the population parameters. Here we evaluated whether different types of reforestations affect Trypoxylon (Trypargilum) lactitarse Saussure (Hymenoptera: Crabronidae) in terms of abundance, percentage of emergence, proportion of males, fluctuating asymmetry, and foraging capacity. We compared primary forest (control) data to data collected in five different habitats: pasture, secondary forest, and tree plantations of Teak. Ficus, and a mixture of native species. The abundance of T. lactitarse was higher in tree plantations than in pasture. However, among the analyzed parameters, Teak plantation presented lower emergence percentage and the majority of individuals born were males. The emerged females in this habitat showed higher asymmetry and lower foraging capacity. Ficus showed lower individual abundance and mixed plantation showed lower emergence percentage, with both plantation types showing higher male emergence. On the other hand, in secondary forest, the analyzed parameters did not differ in relation to the primary forest, being the habitat more efficient in relation to those with tree planting. The changes in population parameters of T. lactitarse in different reforestations and particularly on Teak monocultures were probably attributed to indirect effects, such as low food availability and inadequate environmental conditions.

> services which are important for the maintenance and stability of the ecosystem, leading to enormous ecological, economic, and social losses (Didham 1997, Fontaine *et al* 2006, Barral *et al* 2015, Mitchell *et al* 2015, Fearnside & Figueiredo 2017). Therefore, efforts are increasingly required to reduce the effects of deforestation and to promote biodiversity recovery (Padula & Silva 2005). In fact, every year, billions of dollars are spent on the conservation and recovery of biodiversity (Bueno *et al* 2013). Restoration of an environment is

an expensive intervention, and species chosen for planting in reforested areas can drive the whole ecological succession, influencing regional diversity, ecological interactions, and ecosystem services (Falcão *et al* 2015).

To evaluate the efficiency of these new reforested areas in the recovery of environmental quality, the use of species as bioindicators has proved to be a very effective tool (Edge 2005, Uys *et al* 2006, Engelbrecht 2010, McGeoch *et al* 2011). The sensitivity of some of these species to differences in resources and conditions among different habitats may prevent their occurrence (Dutra & Marco 2015). However, some of these species may colonize and persist on suboptimal habitats, even though suffering indirect effects on their populations. For these species, it is expected that the stress caused by suboptimal conditions may cause changes at the individual and population levels (Clarke 1993, Sanseverino & Nessimian 2008).

Environmental conditions and resources affected by disturbances, such as adverse temperatures, competition, and food scarcity, are known to be factors that may cause alterations in the development of the organisms during their ontogeny (Leung & Forbes 1996; Benítez 2013). One measure commonly used to assess the effect of these disturbances is the increase of the asymmetry (measured by fluctuating asymmetry, hereafter FA) that occurs in bilateral organisms (Palmer & Strobeck 1986). The increase of asymmetry reduces the fitness of the affected organisms in different ways depending on the taxon. However, regardless of the taxon, the asymmetry value found in the individuals allows us to assess the levels of stress to which they were subjected during their development (Leung & Forbes 1996; Pinto et al 2012). Thus, FA can be seen as a metric in the assessment of the quality of the environment for the organisms that live on it (Piscart et al 2005; Benítez 2013).

The variations in habitat quality quickly and strongly influence several insect population parameters, when compared to the other taxa, since they have several generations in a short period of time (Didham 1998; Pearce & Venier 2006; Nichols *et al* 2007). Among insects, the best indicator species are those that spend their whole life within the studied habitats (Clarke *et al* 1986; Camillo *et al* 1995). Thus, wasps that nest in pre-existing cavities fit perfectly as a bioindicator group because they depend on the availability of resources in the local environment to feed their offspring (Evans & Eberhard 1970; Tylianakis *et al* 2005). Additionally, this group is easily sampled using trap nests (Coville 1982), a method that ensures that these species actually inhabit the studied region, thus excluding those that are just passing through (Camillo *et al* 1995).

Known as spider predators, *Trypoxylon* (*Trypargilum*) *lactitarse* Saussure, 1867 (Crabronidae), is considered frequent in trap nests (Beyer *et al* 1987, Morato & Campos 2000, Buschini *et al* 2006, Araujo *et al* 2017), being more abundant in forested environments, but also occurring in deforested areas (Buschini & Wolff 2006). This species is widely distributed from southern Argentina to southern Canada (Coville 1981). Adults use floral resources (pollen and nectar) as energy source, contributing to the pollination of various plant species (Bohart & Menke 1976). However, they also are effective as predators of several species of spiders that are consumed as primary food source during the larval development (Camillo & Brescovit 1999; Buschini *et al* 2008).

As in most of the Hymenoptera, these insects present a haplodiploid system, where unfertilized eggs give rise to males and the fertilized ones give rise to females (Crozier & Pamillo 1996). Due to their ability to control the spermatheca, females are able to choose the sex of each offspring at the moment of oviposition (Godfray 1988, Shintarou & Takayoshi 1999). In addition, there is strong evidence that females are able to adjust the sex ratio of their offspring as a function of environmental conditions (Charnov 1982). In Trypoxylon, one of the main characteristics associated with sexual dimorphism is body size, where females are larger than males and require greater amount of food (Buschini 2007). With this, the availability of resources may be a factor that affects the proportion of the sexes within its offspring (Morato & Campos 2000). Like other solitary wasp species, all food necessary for the development of the larva to adult stage is offered at only once (Molumby 1997, Strohm & Linsenmair 1999). Thus, insufficient food supply at the time of building their breeding cells may lead to stress during offspring development, resulting in an increase in FA of their structures (Niemi & McDonald 2004; Sanseverino & Nessimian 2008). In flying animals, this stress can lead to symmetrical differences in their wings which reduce the ability and aerodynamics of the flight, influencing its adaptive value (Swaddle 1996).

In this study, we evaluated the efficiency of different types of reforested habitats in the recovery of important parameters necessary for the maintenance of the *T. lactitarse* population in Southern Amazon. We specifically tested whether (1) the abundance of individuals, (2) emergence percentage, (3) proportion of males, (4) foraging capacity, and (5) fluctuating asymmetry are influenced by the habitat type in the recovery process when compared with primary forest.

Material and Methods

Study area

We conducted the fieldwork at São Nicolau farm (9°48'S and 58°15'W), located in the municipality of Cotriguaçu in the north of Mato Grosso State, Brazil. The farm has 10,000 ha,

of which 7000 ha are primary forest, 500 ha are secondary forests, 1700 ha are different types of reforestation, and 300 ha are pastures used by cattle. Reforestations and secondary forests were both previously pasture before the revegetation, which occurred between 1999 and 2000 (Rodrigues et al 2011). The regional climate is AW type, according to the Köppen classification (warm and humid), with an average annual temperature of 24°C, 85% humidity, and 2300 mm of precipitation (Rodrigues et al 2011). In this study, we collected data in five different habitats: (1) primary forest (PF) (control), defined as a "terra-firme" and closedcanopy forest without the influence of seasonal flooding of larger rivers. The understory is highly biodiverse, relatively open with a canopy height varying between 30 and 40 m with some trees reaching up to 50 m high; (2) secondary forest (SF), natural restoration without the presence of cattle or anthropogenic interference. The understory of secondary forests is extremely dense, with vegetation higher than 5 m; there is a low leaf litter accumulation and light incidence; (3) teak reforestation (TF) in monodominant stands of the exotic Tectona grandis (Verbenaceae), characterized by an open understory, deciduous leaf-fall pattern in the dry season, and deep leaf litter accumulation; (4) ficus reforestation (FR) in monodominant stands of the native Ficus maxima (Moraceae), sourced from local seed provenance, and characterized by a mixed-age understory shrubs, low leaf litter accumulation, and some patches of grasses; (5) mixed reforestation (MP) of planted native tree species, including Tabebuia chrysotricha (Bignoniaceae), T. roseo-alba, T. impetiginosa, Cedrela fissilis (Meliaceae), Cordia alliodora (Boraginaceae), Simarouba amara (Simaroubaceae), Spondias mombin (Anacardiaceae), Hevea brasiliensis (Euphorbiaceae), and Schizolobium amazonicum (Fabaceae), with an understory of low shrubs and grasses, and low leaf litter accumulation; (6) pasture (PT), dominated by planted grasses (not more than 50 cm tall) used as food for cattle, with very high light incidence, almost no woody vegetation and an absence of leaf litter accumulation (Rodrigues et al 2011). Pasture was used in our design as a "starting point" of forest succession under different reforestations.

Sample design

To sample *T. lactitarse* individuals, we used 10 sample units for each habitat, except for primary forest (n = 14; Fig 1). The minimum distance between each sampling unit of the same treatment was 500 m. In each sample unit, we marked five equidistant points at a distance of 50 m and installed blocks of wood up to 1.5 m high. The wooden blocks were composed of 40 holes distributed equally between the diameters 0.8, 1.2, 1.6, and 2.0 mm by 10 cm deep, where we inserted tubes of black paperboard (Camillo *et al* 1995), totaling 320 blocks and 12,800 trap nests. We inspected the traps every 20 days between August 2016 and July 2017, and those occupied were collected and replaced by another trap with the same measurements. In the laboratory, the trap nests brought from the field were placed in test tubes, closed with cotton, and kept in a dark room at temperatures of between 20°C and 25°C until the emergence of adults. Then, we pinned the emerged adults, quantified the number of cells built, identified their sex, and performed the measurements of wing load (to evaluate the foraging capacity) and fluctuating asymmetry. Voucher specimens were deposited in the Invertebrate Collection of the Universidade Federal de Mato Grosso.

Morphometric measurements

To perform the morphometric measurements of T. lactitarse, we randomly choose only one individual of each sex for each nest that emerged adults, totalizing 119 specimens: PF (male = 10; female = 17), SF (male = 10; female = 18), TR (male = 10; female = 12), FR (male = 10; female = 12), MR (male = 10: female = 10). Due to the low number of individuals that emerged in PT (n = 2, Table 1), it was not possible to include this habitat in the morphometric analyses. To evaluate the fluctuating asymmetry (FA) of individuals, we removed the pair of forewings and photographed and measured the length of each wing with the aid of the software Leica Application Suite version 2.0. The length of the forewing measure starts at the beginning of the costal vein and runs up to the most apical point of the wing. We evaluated the foraging capacity of individuals based on the wing load (WL) represented by the body dry mass (DM) divided by the wing's area (WA). This metric is considered a good predictor of winged individuals' flight capacity (Polidori et al 2013). We measured DM with a precision balance of 0.1 mg and the area of the forewing using the software ImageJ version 1.51n. We calculated WL using the formula WL = DM/WA.

Data analysis

We used generalized linear mixed models (GLMM's) to determine if the abundance, emergence percentage, and proportion of males of *T. lactitarse* differ between primary forest (control) and the habitats studied. For this, we used abundance of individuals, emergence percentage, and proportion of males as response variables. As an explanatory variable (fixed effect) of the model, we used the type of habitat. Considering that we performed repeated measures at each habitat, we used sites as a random effect for all models, since each replicate was visited 12 times (Crawley 2007). For GLMM's implementation, we used the glmmTMB package, and the selection of the best model was given by the comparison using Akaike information criterion.



Fig 1 Map showing the spatial arrangement of our 64 sample units established in six different habitats at the São Nicolau farm, municipality of Cotriguaçu, State of Mato Grosso, Brazil.

We used generalized linear models (GLM's) with Gaussian distribution to determine whether FA and WL varied between habitats. As explanatory variables, we used habitats and the sex of the individuals measured. To verify if the sex of the individuals influenced the response variables within each habitat, we also added to the models the interaction between sex and habitat. After performing the GLMs, we performed the planned comparisons analysis in order to determine if the values of FA and WL (response variables) were different between primary forest and each habitat type. The planned comparison performed for FA and WL was performed separately for each sex. All analyses were performed in program R version 3.5.1 (R Core Team 2018).

Results

During the study period, we collected a total of 839 cells (368 nests) of *T. lactitarse* in the 64 sample units.

The number of cells was highly correlated with the number of nests (Pearson's r = 0.95, p < 0.001, n = 64). Among habitats, PT and FR showed lower abundance of individuals in relation to PF (control), (Z = -3.727, p = 0.000, and Z = -2.460, p < 0.01, respectively). For habitats TR, MR, and SF, no significant differences were found (Fig 2).

The emergence percentage of *T. lactitarse* in the TR, MR, and PT was lower than PF (Z = -2.066, p = 0.038, Z = -4.186, p < 0.001 and Z = -2.043, p = 0.041, respectively). The emergence average in PF was 51.94% of the cells, while in TR 40.21% of the cells built had emerged adults. In MR and PT, the emergence percentage was only 28.74% and 20%, respectively. The emergence percentage of SF and FR did not differ in relation to PF, with 61.18% and 50.53% of individuals born, respectively.

In relation to proportion of males, TR presented the highest deviation, 75.8% of the individuals born were males, followed by FR and MR, with 69.5% and 57.8%,

Table 1 Description of the number of cells built, emergence percentage, proportion of males and females of *Trypoxylon (Trypargilum) lactitarse* individuals born in the habitats studied (mean ± standard error). Statistical values represent the results of planned sex ratio comparisons between pasture (PT), teak reforestation (TR), ficus reforestation (FR), mixed reforestation (MR), and secondary forest (SF) with primary forest (PF).

Environments	No. of cells built	Emergence (%)	Males born (%)	Female born (%)
PT	1.0 ± 0.55*	13.33 ± 10.89*		
TR	16.2 ± 3.57	40.20 ± 2.25*	75.81 ± 2.33*	24.19 ± 3.45
FR	6.4 ± 1.99*	50.53 ± 5.91	69.57 ± 4.43*	30.43 ± 6.27
MR	9.9 ± 3.09	28.74 2.07*	57.81 ± 6.45*	42.19 ± 8.96
SF	12.6 ± 3.43	61.18 ± 1.90	49.31 ± 4.85	50.69 ± 4.79
PF	9.4 ± 2.03	51.94 ± 0.51	43.42 ± 5.54	56.58 ± 4.69

*Coefficients are significantly different from zero (p < 0.05).

respectively. The habitats with the lowest proportions of emerged males were SF (49.31%) and PF (43.42%). When comparing the proportion of males between habitats studied and PF, those actively reforested showed significantly higher values (TR: Z = 3.421, p < 0.000; MR: Z = 2.468, p < 0.013; FR: Z = 2.362, p < 0.018) (Fig 3, Table 1). Due to the low number of emergences in PT (n = 2), this habitat was excluded from the analysis.

The FA of the forewings differed among *T. lactitarse* individuals from the different habitats ($F_{5,118} = 7.222$; p = 0.0001). This asymmetric difference was also associated with the sex of the individuals ($F_{1,113} = 4.560$; p = 0.029). In fact, in all habitats studied, the emerged individuals showed a certain degree of asymmetry in the length of their wings. However, the lowest asymmetry was observed in the PF. In the planned comparisons, we could observe that the females born in the TR presented greater asymmetric variation than the females born in PF ($F_{1,64} = 27.589$; p < 0.000). In the case of males, no differences were observed between the different treatments in relation to the PF (Fig 4, Table 2).

T. lactitarse emerged individuals also differed in WL among habitats ($F_{5,118} = 3.455$; p = 0.002). This difference was associated with sex ($F_{1,113} = 37.100$; p < 0.001) and the variation between sexes was different among habitats ($F_{3,112} = 9.935$; p < 0.001). In PF, while males on average presented the lowest WL values, the females showed WL values close to the highest values recorded for all other habitats (Fig 5). This dimorphism in WL was also observed in MR and SF. Although a similar pattern was observed in FR, this dimorphism was less pronounced. Additionally, in TR, the opposite was found, females and males collected in this habitat presented the lowest and highest values of WL, respectively. When compared with PF, these differences were significant (Fig 5, Table 2).

Discussion

Although T. lactitarse is considered a species of great plasticity, potentially occupying different environments (Pérez-Maluf 1993, Buschini et al 2006, Oliveira-Nascimento & Garófalo 2014, Araujo et al 2017), the different types of habitats studied strongly influenced their nesting behavior and parameters that influence the population structure of their next generation. Biological and structural changes occurring in altered environments may have direct effects on resident species (Morato 2001; Abrahamczyk et al 2011). In the present study, we showed that T. lactitarse occurs and builds nests in all habitat types studied. In all types of tree plantations, T. lactitarse abundance was higher than that in pasture. However, the highest abundance of T. lactitarse occurred in primary forests, probably due to the greater availability of resources and less variation in the environmental conditions in this environment. The lack of vegetation cover and a microenvironment that controls the



Fig 2 Mean number of cells built by *Trypoxylon (Trypargilum) lactitarse* among habitats studied in the southeast of the Amazon: pasture (PT), teak reforestation (TR), ficus reforestation (FR), mixed reforestation (MR), secondary forest (SF), and primary forest (PF).

variation and amplitude in daily temperature was possibly the cause of the low occurrence and high mortality of *T. lactitarse* in pastures. These characteristics are possibly ensured by the



Fig 3 Sex ratio of individuals of *Trypoxylon* (*Trypargilum*) *lactitarse* born in the habitats studied in the southeast of the Amazon: teak reforestation (TR), ficus reforestation (FR), mixed reforestation (MR), secondary forest (SF), and primary forest (PF).



Fig 4 Asymmetric variation in the length of the posterior wings of males and females of *Trypoxylon* (*Trypargilum*) *lactitarse* among habitats studied in the southeast of the Amazon: teak reforestation (TR), ficus reforestation (FR), mixed reforestation (MR), secondary forest (SF), and primary forest (PF).

intensive cattle grazing that hinders the establishment of other species of plants other than grasses (Kurn *et al* 1994, Klein *et al* 2002, Dunne *et al* 2011, Tscharntke *et al* 1998). The species preference for forested environments has also been observed in other studies (Buschini & Wolff 2006; Araujo *et al* 2017).

Although considered generalist, *T. lactitarse* appears to be more flexible in areas with higher prey availability (Buschini *et al* 2008). Spiders commonly predated by *Trypoxylon* usually live on plants where they hunt and breed (Uetz *et al*



Treatments

Fig 5 Variation in wing load of *Trypoxylon* (*Trypargilum*) *lactitarse* males and females emerged in the habitats studied in the southeast of the Amazon: teak reforestation (TR), ficus reforestation (FR), mixed reforestation (MR), secondary forest (SF), and primary forest (PF).

1999, Buschini *et al* 2010, Pitilin *et al* 2012). So, the prey scarcity may also contribute to the low colonization success in the pasture. In studies with other solitary Hymenoptera, spider-hunting wasps have been positively correlated with the complexity of the studied environments (Klein *et al* 2002, Loyola & Martins 2008, Steffan-Dewenter & Tscharntke 2002, Araujo *et al* 2017). This is possibly related to their low dispersal capacity, and due to their specialization in food types, nesting sites, and materials requested for

Table 2 Description of the asymmetry in wing length and wing load of *Trypoxylon (Trypargilum) lactitarse* individuals born in the habitats studied (mean ± standard error). The statistical values represent the results of the planned comparison test of these variables for the male and female individuals born in reforestation in relation to the primary forest.

10

Environments	Sex	Wing asymmetry (mm)			Wing load (mg/mm²)		
		Mean ± SE	F value	p value	Mean ± SE	F value	<i>p</i> value
TR	Female	0.44 ± 0.06	27.59	0.0001*	5.3 ± 0.33	13.25	0.0001*
	Male	0.23 ± 0.09	2.892	0.0959	5.6 ± 0.32	16.57	0.0001*
FR	Female	0.29 ± 0.03	0.273	0.6032	5.5 ± 0.21	2.45	0.1214
	Male	0.22 ± 0.04	0.648	0.425	5.0 ± 0.23	0.653	0.4232
MR	Female	0.34 ± 0.10	3.461	0.0674	5.7 ± 0.20	1.083	0.3018
	Male	0.21 ± 0.04	0.949	0.3353	4.7 ± 0.12	0.64	0.4279
SF	Female	0.18 ± 0.02	1.872	0.176	6.3 ± 0.13	0.373	0.5432
	Male	0.15 ± 0.03	0.705	0.4055	4.8 ± 0.16	0.653	0.4232
PF	Female	0.08 ± 0.01	-	_	6.2 ± 0.15	_	-
	Male	0.09 ± 0.01	-	-	4.3 ± 0.12	-	-

*Coefficients are significantly different from zero (p < 0.05).

Sex

Fema Male nesting (Gathman & Tscharntke 2002, Steffan-Dewenter & Tscharntke 2002, Buschini *et al* 2008, Zurbuchen *et al* 2010). This suggests that all tree plantation types at least favored the colonization of *T. lactitarse* at local scale and, at the same time, the transformation of forests into structurally simple environments such as pastures may lead to its local extinction.

Even though TR presented high T. lactitarse nesting rates in trap nests in relation to other reforestation types, TR did not prove to be adequate for the recovery of the species population patterns observed in PF. This habitat type presented greater mortality of immatures and malebiased sex ratio for individuals born. At the same time, although FR and MR did not show difference in birth rates in relation to PF, they also showed male-biased sex ratio for the emerging offspring. The sex ratio deviation for Trypoxylon species has also been recorded in other studies (Assis & Camillo 1997, Shintarou & Takayoshi 1999, Buschini 2007, Oliveira-Nascimento & Garófalo 2014, Buschini & Bergamaschi 2014). This feature has important consequences for individual fitness and population persistence and may occur in response to variations in environmental conditions, such as low food availability (Hamilton 1979, Molumby 1997, Shintarou & Takayoshi 1999). The sex-biased offspring during the nesting process may occur when there is a strong size dimorphism between sexes (Fisher 1958). Among Trypoxylon, males are noticeably smaller and thus require less investment in food resources for their full development (Buschini et al 2010). Thus, although tree plantations favored the local T. lactitarse occurrence in relation to pasture, a suboptimal availability of food resources may have led the females to invest in a male-biased offspring (Krombein 1967, Sheldon et al 1998, Polidori et al 2013). Another explanation for the smaller number of females in these environments is the difference in mortality between the sexes (Brockmann & Grafen 1992). Since the resource used by one individual is not available to the other (Buschini et al 2010), the lack of sufficient food for the female offspring may have increased its mortality in these environments. As this is a pattern observed at local scale (site), the low availability of females has a negative impact at the population level, reducing mating and nesting rate. This reduces the persistence of T. lactitarse in the planted forests (Hardy et al 1998, 2000, Nunney & Luck 1988). Alternatively, the local persistence of species can be dependent on constant immigration of female individuals from like primary forest.

Among the habitats studied, the individuals of *T. lactitarse* emerged on SF presented morphological patterns more similar to the ones emerged in PF. In tropical environments, forest recovery in abandoned agricultural land can be very fast in the case of low-intensity land use (Chazdon 2003). Soil fertility and the presence of nearby forest remnants are also

factors that facilitate and accelerate the recovery of these environments (Guariguata & Ostertag 2001). In the Western Amazon region, pastures with less than 8 years of abandonment already had tree species growing (Uhl *et al* 1988). All the habitat types of the present study are surrounded by an extensive area of primary Amazon rainforest. Such vast source of propagules should decrease the restoration time on abandoned pastures, creating habitats richer in resources to *T. lactitarse* than reforested sites.

Although studies suggest that in the haplodiploid species of Hymenoptera the diploidism is able to reduce the effects of the stress in function of heterozygosity (Clarke et al 1986, Tomkins & Kotiaho 2001, Miklasevskaja & Packer 2015), our study shows that females (diploids) were more asymmetric than males. In addition, the variation in asymmetry of females emerged in TR was higher in relation to the ones emerged in PF. Curiously, males show no significant variation between habitats. Environments with some type of restriction may impose difficulties to the establishment of colonizing species, promoting low heterozygosity due the high rates of inbreeding (Mitton & Grant 1984). Therefore, individuals and populations with low heterozygosity are sensitive to disturbances during their development (Clarke 1993; Tomkins & Kotiaho 2001). This suggests that although T. lactitarse females are able to nest in TR, this habitat may exert pressures on the quality of their offspring. So, synergistically with the male-biased populations, the habitat can also negatively affect the quality of females born. We believe that these values were less accentuated for males by the fact that they needed fewer resources for their development. Thus, in a resource-deficient environment, females are more susceptible to development problems than males, which may account for the greater male birth in habitats like TR, MR, and FR.

The effect of environmental stress on the FA increase of organisms has been already reported in several groups, such as other insects (Chang et al 2007, Pinto et al 2012, Banaszak-Cibicka et al 2018), birds (Vangestel et al 2011), crustaceans (Ho et al 2009), fishes (Vandenbussche et al 2018), and plants (Lobregat et al 2017). The asymmetric differences, as observed between the wings of the T. lactitarse females in TR, potentially affect their adaptive success (Swaddle 1996). The disparity in these structures decreases the capability to escape from predators by reducing the efficiency in take-off and maneuverability (Tomkins & Kotiaho 2001), as well as reducing foraging abilities (Swaddle et al 1997; Samejima & Tsubaki 2010). A study on parasitic wasps (Bennett & Hoffmann 1998) showed that the asymmetry of the wings was negatively associated with the ability to find the eggs of the host (Lepidoptera). In addition, the impairment of foraging abilities may affect the female's fitness during the nesting process (Thomas 1993). Therefore, even though adults build nests with relatively high abundance, the females of

T. lactitarse populations born in TR are particularly more asymmetrical, decreasing the potential of population persistence on such habitats.

Females born in TR also differed in WL relative to the females emerged in PF. A lower WL reflects greater flight efficiency, since they have larger wings in relation to the body (Polidori et al 2013). This feature may have favored the success in the occupation of this reforestation by the founding females. We just measured the WL of the females born in TR, not the adult ones. However, if WL is genetically determined (Endler 1977, Lack et al 2015), the population of T. lactitarse is composed by females that behave as longdistance dispersers and it reinforces the hypothesis of the dependence of the populations living in TR on females constantly migrating from other sites. Additionally, large wings need more energy for their development and, in an environment with limiting resources, these structures are more susceptible to the FA caused by stress during their development (Leung & Forbes 1996). Although larger wings may be a feature that facilitated the occupation in TR by females, it possibly does not remain stable throughout the generations. The males born in TR have an inverse WL pattern in relation to females. Their higher WL reflects a higher body weight (Polidori et al 2013). Studies indicate that the increase in WL is positively related to the increase in the flying musculature (Kemp & Alcock 2008; Yao 2011). Although greater flight musculature results in higher energy expenditure, individuals with these characteristics present greater maneuverability and acceleration during their foraging (Marden 1989). In a study on a "cicada killer" wasp, greater competitive success was observed in territorial males with higher flying musculature (Coelho & Holliday 2001), and the same pattern was found for dragonflies (Marden 1989). Some studies show that Trypoxylon males may exhibit territorial behavior to increase the chances of reproductive success (Alcock 1975; Buschini & Donatti 2012); therefore, a greater flight musculature may give them more success for intercepting the longdistance female since they can take off with greater speed (Byrne et al 1988). Thus, environments with low availability of females may have favored males with higher WL, since this feature seems to increase the success in interception of females for mating.

As expected, among the different habitat types evaluated, pastures were inadequate for the establishment of *T. lactitarse*. This demonstrates the need for reforestation to recover local populations from deforested areas. However, the different types of tree plantations presented a lower efficiency in the recovery of population patterns in relation to natural regeneration areas (secondary forest). The planted species used in the present study, especially *Tectona grandis*, probably did not favor the recovery of environmental conditions that could aid in the re-establishment of *T. lactitarse* population patterns. The success of secondary

forest was possibly due to its proximity to the primary forest (source of colonizing species). This may have facilitated the establishment of native species and consequently led to the development of more similar characteristics found in natural environments, such as food availability and adequate environmental conditions. However, the effectiveness of secondary forests from abandoned areas may fail in regions without the presence of nearby natural forests. This demonstrates that environmental recovery projects should be evaluated individually, taking into account the characteristics of each area to be recovered.

Acknowledgments We highly acknowledge the Office National des Forêts of Brazil for the logistical support and Coordenação de Aperfeiçoamento de Pessoal de Nível Superior for the scholarships provided to GJA.

Author Contributions GJA, DST and TJI Conceptualization, GJA, DST and TJI Data curation, GJA, DST and TJI Formal analysis, TJI Funding acquisition, GJA, DST and TJI Investigation, GJA and TJI Methodology, TJI Project administration, TJI Resources, TJI Supervision, GJA, DST and TJI Validation, GJA, DST and TJI Visualization, GJA and TJI Writing (original draft preparation), GJA, DST and TJI Writing (review and editing).

References

- Abrahamczyk S, Gottleuber P, Matauschek C, Kessler M (2011) Diversity and community composition of euglossine bee assemblages (Hymenoptera: Apidae) in western Amazonia. Biodivers Conserv 20: 2981–3001
- Alcock J (1975) Social interactions in the solitary wasp, Cerceris simplex (Hymenoptera: Sphecidae). Behavior 54:142–152
- Araujo GJ, Fagundes R, Antonini Y (2017) Trap-nesting hymenoptera and their network with parasites in recovered Riparian forests Brazil. Neo Entomol 47:26–36. https://doi.org/10.1007/s13744-017-0504-4
- Assis JMF, Camillo E (1997) Diversidade, sazonalidade e aspectos biológicos de vespas solitárias (Hymenoptera: Sphecidae: Vespidae) em ninhos armadilhas na região de Ituiutaba. An Soc Entomol Bras 26: 335–347
- Banaszak-Cibicka W, Fliszkiewicz M, Langowska ZM (2018) Body size and wing asymmetry in bees along an urbanization gradient. Apidologie 49:297–306
- Barral MP, Benayas JMR, Meli P, Maceira NO (2015) Quantifying the impacts of ecological restoration on biodiversity and ecosystem services in agroecosystems: a global meta-analysis. Agri Eco Envirol 202: 223–231
- Benítez HA (2013) Assessment of patterns of fluctuating asymmetry and sexual dimorphism in carabid body shape. Neo Entomol 42: 164–169
- Bennett DM, Hoffmann AA (1998) Effects of size and fluctuating asymmetry on field fitness of the parasitoid Trichogramma carverae (Hymenoptera: Trichogrammatidae). J Anim Ecol 67:580–591
- Beyer WN, Miller GW, Fleming WJ (1987) Populations of trap-nests wasps near a major source of fluoride emissions in Western Tennessee. Proc Entomol Soc Wash 89:478–482

- Bohart RM, Menke AS (1976) Sphecid wasps of the world a generic revision. University of California Press, London, p 660
- Brockmann HJ, Grafen A (1992) Sex rations and life-history patterns of a solitary wasp, Trypoxylon (Trypargilum) politum (Hymenoptera: Sphecidae). Behav Ecol Sociobiol 30:7–27
- Bueno AF, Paula-Moraes SV, Gazzoni DL, Pomari AF (2013) Economic thresholds in soybean-integrated pest management: old concepts, current adoption, and adequacy. Neo Entomol 42:439–447
- Buschini MLT (2007) Life-history and sex allocation in *Trypoxylon (syn. Trypargilum) lactitarse* (Hymenoptera; Crabronidae). J Zool Syst Evol Res 45:206–213
- Buschini MLT, Bergamaschi ABC (2014) Sex ratio and parental investment in *Trypoxylon* (*Trypargilum*) agamemnon Richards (Hymenoptera, Crabronidae). Braz J Biol 74:231–237
- Buschini MLT, Donatti AJ (2012) Nesting behavior of *Trypoxylon* (*Trypargilum*) agamemnon Richards (Hymenoptera: Crabronidae). Braz J Biol 72:353–362
- Buschini MLT, Wolff LL (2006) Notes on the biology of *Trypoxylon* (*Trypargilum*) opacum Brethes (Hymenoptera: Crabronidae) in southern Brazil. Braz J Biol 66:915–926
- Buschini MLT, Lazzarini LW, Niesting F (2006) Nesting biology of *Trypoxylon (Trypargilum) lactitarse* (Hymenoptera; Crabronidae) in trap-nests in southern Brazil. Braz J Biol 66:161–171
- Buschini MLT, Borda NA, Brescovit AD (2008) Patterns of prey selection of *Trypoxylon (Trypargilum) lactitarse* Sausurre (Hymenoptera: Crabronidae) in southern Brazil. Braz J Biol 68:519–528
- Buschini MLT, Caldas TR, Borba NA, Brescovit AD (2010) Spiders used as prey by the hunting wasp *Trypoxylon (Trypargilum) agamemnon* Richards (Hymenoptera: Crabronidae). Zool Stud 49:169–175
- Byrne DN, Buchmann SL, Spangler HG (1988) Relationship between wing loading, wingbeat frequency and body mass in homopterous insects. J Exp Biol 135:9–24
- Camillo E, Brescovit AD (1999) Aspectos biológicos de *Trypoxylon* (*Trypargilum*) *lactitarse* Saussure e *Trypoxylon* (*Trypargilum*) *rogenhoferi* Kohl (Hymenoptera: Sphecidae) em ninhos-armadilhas, com especial referência a suas presas. An Soc Entomol Bras 28:251– 262
- Camillo E, Garofalo CA, Serrano JC, Muccillo G (1995) Diversidade e abundância sazonal de abelhas e vespas solitárias em ninhos armadilhas (Hymenoptera: Apocrita: Aculeata). Rev Bras Entomol 39:459– 470
- Chang CH, Lin YH, Chen IH, Chuang SC, Chen JH (2007) Taxonomic reevaluation of the Taiwanese montane earthworm Amynthas wulinensis Tsai, Shen & Tsai, 2001 (Oligochaeta: Megascolecidae): polytypic species or species complex? Org Divers Evol 7:231–240
- Charnov EL (1982) The theory of sex allocation. Princeton University Press, Princeton, p 355
- Chazdon RL (2003) Tropical forest recovery: legacies of human impact and natural disturbances. Perspect Plant Ecol Evol Syst 6:51– 71
- Clarke GM (1993) Patterns of developmental stability of *Chrysopa perla L*. (Neuroptera: Chrysopidae) in response to environmental pollution. Environ Entomol 22:1362–1366
- Clarke GM, Brand GW, Whitten MJ (1986) Fluctuating asymmetry: a technique for measuring developmental stress caused by inbreeding. Aust J Biol Sci 39:145–153
- Coelho JR, Holliday CW (2001) Effects of size and flight performance on intermale mate competition in the cicada killer, *Sphecius speciosus* (Hymenoptera: Sphecidae). J Insect Behav 14:345–351
- Coville RE (1981) Biological observations on three *Trypoxylon* wasps in the subgenus *Trypargilum* from Costa Rica: *T. nitidum schultessi*, *T. saussurei* and *T. lactitarse* (Hymenoptera: Sphecidae). Pan Pac Entomol 57:332–340
- Coville RE (1982) Wasps of the genus *Trypoxylon* subgenus *Trypargilum* in North America. University of California Press, Berkeley, p 163

- Crawley MJ (2007) The R book. Imperial College London, Silwood Park Chichester, p 1080
- Crozier RH, Pamillo P (1996) One into two will go. Nature 383:574–575
- Didham RK (1997) The infuence of edge effects and forest fragmentation on leaf-litter invertebrates in Central Amazonia. In: Laurance WF, Bierregaard RO (eds) Tropical forest remnants: ecology, Manag Cons Frag Communt Chicago, pp 55–70
- Didham RK (1998) Altered leaf-litter decomposition rates in tropical forest fragments. Oecologia 116:397–406
- Dunne T, Western D, Dietrich WE (2011) Effects of cattle trampling on vegetation, infiltration, and erosion in a tropical rangeland. J Arid Environ 75:58–69
- Dutra S, Marco P (2015) Bionomic differences in odonates and their influence on the efficiency of indicator species of environmental quality. Ecol Indic 49:132–142
- Edge DA (2005) Butterfly conservation in the southern Cape. Metamorphosis 16:28–46
- Endler JA (1977) Geographic variation, speciation and clines. Monogr Popul Biol 10:1–246
- Engelbrecht IA (2010) Invertebrate species inventories in protected area management: are they useful? Afr Entomol 18:235–245
- Evans HE, Eberhard MJ (1970) The wasps. The University of Michigan Press, Ann Arbor, p 265
- Falcão CF, Dátillo W, Izzo TJ (2015) Efficiency of different planted forests in recovering biodiversity and ecological interactions in Brazilian Amazon Forest. For Ecol Manag 339:105–111
- Fearnside PM (2006) Deforestation in Amazonia: dynamics, impacts and control. Acta Amaz 36:395–400
- Fearnside P, Figueiredo A (2017) China's influence on deforestation in Brazilian Amazonia: a growing force in the State of Mato Grosso. In: Ray R, Gallagher K, Lopez A, Sanborn C (eds) China and sustainable development in Latin America: the social and environmental dimension, London, pp 229–266
- Fisher RA (1958) The genetical theory of natural selection. The Clarendon Press, New York, p 308
- Fontaine C, Dajoz I, Meriguet J, Loreau M (2006) Functional diversity of plant–pollinator interaction webs enhances the persistence of plant communities. PLoS Biol 4:129–135
- Gathman A, Tscharntke T (2002) Foraging ranges of solitary bees. J Anim Ecol 71:757–764
- Gibbs HK, Ruesch AS, Achard F, Clayton MK, Holmgren P, Ramankutty N, Foley JA (2010) Tropical forests were the primary sources of new agricultural land in the 1980s and 1990s. PNAS 107:16732–16737
- Godfray HCJ (1988) Virginity in haplodiploid populations: a study on lig wasps. Eco Entomol 13:283–291
- Guariguata MR, Ostertag R (2001) Neotropical secondary forest succession: changes in structural and functional characteristics. For Ecol Manag 148:185–206
- Hamilton WD (1979) Wingless and fighting males in fig wasps and other insects. In: Blum MS, Blum NA (eds) Reproductive competition and sexual selection in insects. Academic Press, London, pp 167–220
- Hardy ICW, Dijkstra LJ, Gillis JEM, Luft PA (1998) Patterns of sex ratio, virginity and developmental mortality in gregarious parasitoids. Biol J Linn Soc 64:239–270
- Hardy ICW, Stokkebo S, Bùnlùkke-Pedersen J, Sejr MK (2000) Insemination capacity and dispersal in relation to sex allocation decisions in Goniozus legneri (Hymenoptera: Bethylidae): why are there more males in larger broods? Ethology 106:1021–1032
- Ho GWC, Leung KMY, Lajus D, Ng JSS, Chan BKK (2009) Fluctuating asymmetry of Amphibalanus (Balanus) amphitrite (Cirripedia: Thoracica) in association with shore height and metal pollution. Hydrobiologia 621:21–32
- Kemp DJ, Alcock J (2008) Aerial contests, sexual selection and flight morphology in solitary Pompilid wasps. Ethology 114:195–202

- Klein AM, Steffan-Dewenter I, Tscharntke T (2002) Effects of land-use intensity in tropical agroforestry systems on flower-visiting and trapnesting bees and wasps. Conserv Biol 16:1003–1014
- Krombein KV (1967) Trap-nesting wasps and bees: life histories, nests and associates. Washington, p 569
- Kurn D, Bretz S, Akbari H (1994) The potential for reducing urban air temperatures and energy consumption through vegetative cooling. ACEEE summer study on Energy Efficiency in Buildings, America Council for an Energy Efficient Economy. Pacific Grove: American Council for an Energy Efficient Economy. http://www.osti.gov/ bridge/servlets/purl/10180633-hLSlld/native/10180633.pdf. Accessed 11 August 2017
- Lack JB, Monette MJ, Johanning EJ, Sprengelmeyer QD, Pool JE (2015) Decanalization of wing development accompanied the evolution of large wings in high-altitude Drosophila. PNAS 113:1014–1019
- Laurance WF, Camargo JLC, Luizão RCC, Laurance SG, Pimm SL, Bruna EM, Stouffer PC, Bruce Williamson G, Benítez-Malvido J, Vasconcelos HL, van Houtan KS, Zartman CE, Boyle SA, Didham RK, Andrade A, Lovejoy TE (2011) The fate of Amazonian forest fragments: a 32-year investigation. Bio Conserv 144:56–67
- Leung B, Forbes MR (1996) Fluctuating asymmetry in relation to stress and fitness: effects of trait type as revealed by meta-analysis. Ecoscience 3:400–413
- Lobregat G, Perilli MLL, Neves F, Campos R (2017) Fluctuating asymmetry, leaf thickness and herbivory in Tibouchina granulosa: an altitudinal gradient analysis. Arthropod Plant Interact 12:277–282. https:// doi.org/10.1007/s11829-017-9568-7
- Loyola RD, Martins RP (2008) Habitat structure components are effective predictors of trap-nesting Hymenoptera diversity. Basic Appl Ecol 9:765–742
- Marden JH (1989) Body building dragonflies: costs and benefits of maximizing flight muscle. Physiol Zool 62:505–521
- McGeoch MA, Sithole H, Samways MJ, Simaika JP, Pryke JS, Picker M, Uys C, Armstrong AJ, Dippenaar-Schoeman AS, Engelbrecht IA, Braschler B, Hamer M (2011) Conservation and monitoring of invertebrates in terrestrial protected areas. Koedoe 53:1–13
- Miklasevskaja M, Packer L (2015) Fluctuating asymmetry in am extreme morphological adaptation in the Chilean bee *Xeromelissa rozeni* (Hymenoptera: Colletidae). Can J Zool 93:833–840
- Mitchell MG, Suarez-Castro AF, Martinez-Harms M, Maron M, McAlpine C, Gaston KJ, Johansen K, Rhodes JR (2015) Reframing landscape fragmentation's effects on ecosystem services. Trends Ecol Evol 30:190– 198
- Mitton JB, Grant MC (1984) Association among protein heterozygosity, growth rate and developmental homeostasis. Ann Rev Ecol System 15:479–499
- Molumby A (1997) Why make daughter larger? Maternal sex-allocation and sex-dependent selection for body size in a mass-provisioning wasp *Trypoxylon politum*. Behav Ecol 8:279–287
- Morato EF (2001) Biologia e ecologia de Anthodioctes moratoi Urban (Hymenoptera, Megachilidae, Anthidiini) em matas contínuas e fragmentos na Amazônia Central, Brasil. Rev Bras Zool 18:729–736
- Morato EF, Campos LAO (2000) Efeitos da fragmentação florestal sobre vespas e abelhas solitárias em uma área da Amazônia Central. Rev Bras Zool 17:429–444
- Newbold T, Hudson LN, Hill SLL, Contu S, Lysenko I, Senior RA, Börger L, Bennett DJ, Choimes A, Collen B, Day J, de Palma A, Díaz S, Echeverria-Londoño S, Edgar MJ, Feldman A, Garon M, Harrison MLK, Alhusseini T, Ingram DJ, Itescu Y, Kattge J, Kemp V, Kirkpatrick L, Kleyer M, Correia DLP, Martin CD, Meiri S, Novosolov M, Pan Y, Phillips HRP, Purves DW, Robinson A, Simpson J, Tuck SL, Weiher E, White HJ, Ewers RM, Mace GM, Scharlemann JPW, Purvis A (2015) Global effects of land use on local terrestrial biodiversity. Nature 520:45–50
- Nichols E, Larsen T, Spector S, Davis AL, Escobar F, Favila M, Vulinec K (2007) Global dung beetle response to tropical forest modification

and fragmentation: a quantitative literature review and meta-analysis. Bio Conserv 137:1–19

- Niemi GJ, McDonald ME (2004) Application of ecological indicators. An Rev Ecol Evol Syst 35:89–111
- Nunney L, Luck RF (1988) Factors influencing the optimum sex ratio in structured populations. J Theor Biol 33:1–30
- Oliveira-Nascimento AL, Garófalo CA (2014) Trap-nesting solitary wasps (Hymenoptera: Aculeata) in an insular landscape: mortality rates for immature wasps, parasitism, and sex ratios. Sociobiology 61:207–217
- Padula RC, Silva LP (2005) Gestão e licenciamento ambiental no Brasil: modelo de gestão focado na qualidade do meio ambiente. Cadernos EBAPE.BR, Rio de Janeiro, pp 1–15
- Palmer AR, Strobeck C (1986) Fluctuating asymmetry: measurement, analysis, patterns. Annu Rev Ecol Syst 17:391–421
- Pearce JL, Venier LA (2006) The use of ground beetles (Coleoptera: Carabidae) and spiders (Araneae) as bioinidcators of sustainable forest management: a review. Ecol Indic 6:780–793
- Pérez-Maluf R (1993) Biologia de vespas e abelhas solitárias, em ninhos armadilhas em Viçosa - MG. Dissertation, Universidade Federal de Viçosa
- Pinto NS, Juen L, Cabette HSR, Júnior PM (2012) Fluctuating asymmetry and wing size of Argia tinctipennis Selys (Zygoptera: Coenagrionidae) in relation to riparian forest preservation status. Neo Entomol 41:178– 185
- Piscart C, Moreteau JC, Beisel JN (2005) Biodiversity and structure of macroinvertebrate communities along a small permanent salinity gradient (Meurthe River, France). Hydrobiologia 551:227–236
- Pitilin RB, Araújo MS, Buschini MLT (2012) Individual specialization in the hunting-wasp *Trypoxylon (Trypargilum) agamemnon* Richards (Hymenoptera: Crabronidae). Zool Stud 51:655–662
- Polidori C, Santoro D, Bluthgen N (2013) Does prey mobility affect niche width and individual specialization in hunting wasps? A networkbased analysis. Oikos 122:385–394
- R Development Core Team R (2018) A language and environment for statistical computing. Version 3.5.1 [software]. User's guide and application published http://www.R-project.org
- Rodrigues DJ, Izzo TJ, Battirola LD (2011) Descobrindo a Amazônia Meridional: Biodiversidade da Fazenda São Nicolau. Pau e Prosa Comunicação, Cuiabá, p 301
- Samejima Y, Tsubaki Y (2010) Body temperature and body size affect flight performance in a damselfly. Behav Ecol Sociobiol 64:685– 692
- Sanseverino AM, Nessimian JL (2008) Larvas de Chironomidae (Diptera) em depósitos de folhiço submerço em um richo de primeira ordem da Mata Atlântica (Rio de Janeiro, Brasil). Rev Bras Entom 52:95–104
- Sheldon BC, Merilä J, Lindgren G, Ellegren H (1998) Gender and environmental sensitivity in nestling collared flycatchers. Ecol press 79:1939– 1948
- Shintarou O, Takayoshi N (1999) Factors affecting female-biased sex ratio in a trap-nesting wasp, *Trypoxylon malaisei*. Rev Popul Ecol 41: 169–175
- Steffan-Dewenter I, Tscharntke T (2002) Insect communities and biotic interactions on fragmented calcareous grassland: a mini review. Biol Conserv 104:275–284
- Strohm E, Linsenmair KE (1999) Measurement of parental investment and sex allocation in the European beewolf *Philanthus triangulum* F. (Hymenoptera: Sphecidae). Behav Ecol Sociobiol 47:76–88
- Swaddle JP (1996) Reproductive success and symmetry in zebra finches. Anim Behav 51:203–210
- Swaddle JP, Witter MS, Cuthill A, McCowen P (1997) Plunage condition affects flight performance in starlings: implications for developmental homeostasis, abrasion and moult. J Avian Biol 27:103–111
- Thomas ALR (1993) On the aerodynamics of birds' tails. Philos Trans R Soc Lond B 340:361–380
- Tomkins JL, Kotiaho JS (2001) Fluctuating asymmetry. Encyclopedia of life and sciences. https://doi.org/10.1038/npg.els.0003741

- Tscharntke T, Gathmann A, Steffandewenter I (1998) Bioindication using trap-nesting bees and wasps and their natural enemies: community structure and interactions. J Appl Ecol 35:708–719
- Tylianakis JM, Klein AM, Tscharntke T (2005) Spatiotemporal variation in the effects of a tropical habitat gradient on Hymenoptera diversity. Ecology 86:3296–3302
- Uetz GW, Halaj J, Cady AB (1999) Guild structure of spiders in major crops. J Arachnol 27:270–280
- Uhl C, Clark K, Maquirino P (1988) Vegetation dynamics in Amazonian treefall gaps. Ecology 69:751–763
- Uys C, Hamer M, Slotow R (2006) The effect of burn area on invertebrate recolonisation in grasslands in the Drakensberg, South Africa. Afr Zool 41:51–65
- Vandenbussche PSP, Spennato G, Pierson PM (2018) Assessment of the use of Oblada melanura (L. 1758) otolith fluctuating asymmetry as environmental disturbance indicator. Mar Environ Res 136:48–53

- Vangestel C, Braeckman BP, Matheve H, Lens L (2011) Constraints on home range behaviour affect nutritional condition in urban house sparrows (*Passer domesticus*). Biol J Linn Soc 101:41–50. https://doi. org/10.1111/j.1095-8312.2010.01493.x
- Yao I (2011) Phylogenetic comparative methods reveal higher wing loading in ant-attended Tuberculatus aphids (Hemiptera: Aphididae). Can Entomol 143:34–43
- Zurbuchen A, Bachofen C, Müller A, Hein S, Dorn S (2010) Are landscape structures insurmountable barriers for foraging bees? A markrecapture study with two solitary pollen specialist species. Apidologie 41:497–508

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.